ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 72 • Numbers 1–4 • 2005

Author index

AHMED, M.

Some observations on ticks (Acari: Ixodidae) infesting sheep in River Nile Province of Northern Sudan—M. AHMED, A.M. EL HUSSEIN & A.O. EL KHIDER 239–243

ALBERTYN, J.

Psittacine beak and feather disease virus haemagglutinating activity using erythrocytes from African Grey parrots and Brown-headed parrots—K. KONDIAH, J. ALBERTYN & R.R. BRAGG 263–265

ALMLI, B.

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

AVENANT-OLDEWAGE, A.

Aspects of the ecology of the Asian tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934 in yellowfish in the Vaal Dam in South Africa—A. BERTASSO & A. AVENANT-OLDEWAGE 207–217

BASTIANELLO, S.S.

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

BENGIS, R.G.

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

BERTASSO, A.

Aspects of the ecology of the Asian tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934 in yellowfish in the Vaal Dam in South Africa—A. BERTASSO & A. AVENANT-OLDEWAGE 207–217

BEZUIDENHOUT, A.J.

A lectin histochemical study of the thoracic respiratory air sacs of the fowl—A.J. BEZUIDENHOUT 175–180

BOSHOFF, H.M.

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRECEK 185–187

BOTHA, C.J.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

BRAACK, L.E.O.

Small mammals as hosts of immature ixodid ticks—I.G. HORAK, L.J. FOURIE & L.E.O. BRAACK 255–261

BRAGG, R.R.

Effects of differences in virulence of different serovars of *Haemophilus paragallinarum* on perceived vaccine efficacy—R.R. BRAGG 1–6

Psittacine beak and feather disease virus haemagglutinating activity using erythrocytes from African Grey parrots and Brown-headed parrots—K. KONDIAH, J. ALBERTYN & R.R. BRAGG 263–265

BREMER, C.W.

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

BROWN, C.G.D.

In vivo comparison of susceptibility between *Bos indicus* and *Bos taurus* cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

CINDZI, Z.M.

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

COETZER, J.A.W.

The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques—E.S.M. TUPPURAINEN, E.H. VENTER & J.A.W. COETZER 153–164

COUACY-HYMANN, E.

The phytochemical composition and *in vitro* antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (*Combretaceae*) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

DE KOCK, K.N.

Distribution, habitats and role as intermediate host of the freshwater snail, *Bulinus forskalii*, in South Africa—K.N. DE KOCK & C.T. WOLMARANS 165–174

DIGIANI, M.-C.

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

DOBORO, F.A.

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

DOLAN, T.T.

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

In vivo comparison of susceptibility between Bos indicus and Bos taurus cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

DUNGU, B.

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

DURAND, A.M.

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

DURAND, D.T.

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRECEK 185–187

DURETTE-DESSET, M.-C.

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

EL HUSSEIN, A.M.

Some observations on ticks (Acari: Ixodidae) infesting sheep in River Nile Province of Northern Sudan—M. AHMED, A.M. EL HUSSEIN & A.O. EL KHIDER 239–243

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

EL KHIDER, A.O.

Some observations on ticks (Acari: Ixodidae) infesting sheep in River Nile Province of Northern Sudan—M. AHMED, A.M. EL HUSSEIN & A.O. EL KHIDER 239–243

ELOFF, J.N.

Is the use of *Gunnera perpensa* extracts in endometritis related to antibacterial activity?—L.J. McGAW, R. GEHRING, L. KATSOULIS & J.N. ELOFF 129–134

EMIKPE, B.O.

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

FLÅØYEN, A.

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

FOURIE, L.J.

Small mammals as hosts of immature ixodid ticks—I.G. HORAK, L.J. FOURIE & L.E.O. BRAACK 255–261

Elephant shrews as hosts of immature ixodid ticks—L.J. FOURIE, I.G. HORAK & P.F. WOODALL 293–301

FOURIE, N.

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

GEHRING, R.

Is the use of *Gunnera perpensa* extracts in endometritis related to antibacterial activity?—L.J. McGAW, R. GEHRING, L. KATSOULIS & J.N. ELOFF 129–134

HASSAN, S.M.

Natural infection rates and transmission of Theileria annulata by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

HLATSHWAYO, M.

A survey of tick control methods used by resource-poor farmers in the Qwa-Qwa area of the eastern Free State Province, South Africa—M. HLATSHWAYO & P.A. MBATI 245–249

HORAK, I.G.

Small mammals as hosts of immature ixodid ticks—I.G. HORAK, L.J. FOURIE & L.E.O. BRAACK 255–261

Elephant shrews as hosts of immature ixodid ticks—L.J. FOURIE, I.G. HORAK & P.F. WOODALL 293–301

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Ticks on crested francolins, *Francolinus sephaena*, and on the vegetation on a farm in Limpopo Province, South Africa—A.C. UYS & I.G. HORAK 339–343

HOVE, T.

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Seroprevalence of *Toxoplasma gondii* infection in domestic pigs reared under different management systems in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 231–237

Seroprevalence of *Toxoplasma gondii* infection in goats and sheep in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 267–272

JAMAL, S.

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

KALEMA-ZIKUSOKA, G.

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

KATSOULIS, L.

Is the use of *Gunnera perpensa* extracts in endometritis related to antibacterial activity?—L.J. McGAW, R. GEHRING, L. KATSOULIS & J.N. ELOFF 129–134

KELLERMAN, T.S.

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

KELLY, P.J.

A preliminary study to evaluate the immune responses induced by immunization of dogs with inactivated *Ehrlichia canis* organisms—S. MAHAN, P.J. KELLY & S.M. MAHAN 119–128

KER, J.

Electrocardiographic surrogates of structural myocardial alterations in the Dorper sheep heart—J. KER & E.C. WEBB 273–277

KIMBITA, E.N.

Immunization of young chicks using graded dose of wild strain of *Eimeria tenella*—E.N. KIMBITA, P.M. SAMANI, E.N. MACHANGE & R.S. SILAYO 251–254

KIRBERGER, R.M.

A canine model of normovolaemic acute anaemia—T.C. SPOTSWOOD, R.M. KIRBERGER, L.M.P.K. KOMA, F. REYERS & L. VAN DER MERWE 135–143

KOMA, L.M.P.K.

A canine model of normovolaemic acute anaemia—T.C. SPOTSWOOD, R.M. KIRBERGER, L.M.P.K. KOMA, F. REYERS & L. VAN DER MERWE 135–143

KONDIAH, K.

Psittacine beak and feather disease virus haemagglutinating activity using erythrocytes from African Grey parrots and Brown-headed parrots—K. KONDIAH, J. ALBERTYN & R.R. BRAGG 263–265

KRECEK, R.C.

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRECEK 185–187

KRÜGER, L.P.

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

LAMIEN, C.E.

The phytochemical composition and *in vitro* antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (*Combretaceae*) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

LAWRENCE, A.N.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

LAZARUS, A.G.

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

LIND, P.

Seroprevalence of *Toxoplasma gondii* infection in domestic pigs reared under different management systems in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 231–237

Seroprevalence of *Toxoplasma gondii* infection in goats and sheep in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 267–272

MACHANGE, E.N.

Immunization of young chicks using graded dose of wild strain of *Eimeria tenella*—E.N. KIMBITA, P.M. SAMANI, E.N. MACHANGE & R.S. SILAYO 251–254

MACUAMULE, C.

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

MAHAN, S.

A preliminary study to evaluate the immune responses induced by immunization of dogs with inactivated *Ehrlichia canis* organisms—S. MAHAN, P.J. KELLY & S.M. MAHAN 119–128

MAHAN, S.M.

A preliminary study to evaluate the immune responses induced by immunization of dogs with inactivated *Ehrlichia canis* organisms—S. MAHAN, P.J. KELLY & S.M. MAHAN 119–128

MAICHOMO, M.W.

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

MAONONGA, D.B.

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

MARCOTTY, T.

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

MATENGA, ELIZABETH

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

MATETE, G.O.

Association of trypanosomosis risk with dairy cattle production in western Kenya—G.L. MUGUNIERI & G.O. MATETE 279–284

MAVENYENGWA, M.

A macro- and light microscopical study of the pathology of *Calicophoron microbothrium* infection in experimentally infected cattle—M. MAVENYENGWA, S. MUKARATIRWA, M. OBWOLO & J. MONRAD 321–332

MBATI, P.A.

A survey of tick control methods used by resource-poor farmers in the Qwa-Qwa area of the eastern Free State Province, South Africa—M. HLATSHWAYO & P.A. MBATI 245–249

MBOGO, S.K.

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

McGAW, L.J.

Is the use of *Gunnera perpensa* extracts in endometritis related to antibacterial activity?—L.J. McGAW, R. GEHRING, L. KATSOULIS & J.N. ELOFF 129–134

MEDA, A.

The phytochemical composition and *in vitro* antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (*Combretaceae*) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

MEINTJES, R.A.

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

MICHAEL, L.M.

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRECEK 185–187

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

MICHEL, A.L.

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

MONRAD, J.

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

A macro- and light microscopical study of the pathology of *Calicophoron microbothrium* infection in experimentally infected cattle—M. MAVENYENGWA, S. MUKARATIRWA, M. OBWOLO & J. MONRAD 321–332

MUGUNIERI, G.L.

Association of trypanosomosis risk with dairy cattle production in western Kenya—G.L. MUGUNIERI & G.O. MATETE 279–284

MUKARATIRWA, S.

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Seroprevalence of *Toxoplasma gondii* infection in domestic pigs reared under different management systems in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 231–237

Seroprevalence of *Toxoplasma gondii* infection in goats and sheep in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 267–272

A macro- and light microscopical study of the pathology of *Calicophoron microbothrium* infection in experimentally infected cattle—M. MAVENYENGWA, S. MUKARATIRWA, M. OBWOLO & J. MONRAD 321–332

MUSONDA, M.M.

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

MUTUGI, J.J.

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

MWASE, M.

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

NACOULMA, O.G.

The phytochemical composition and *in vitro* antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (*Combretaceae*) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

NAUDÈ, T.W.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÈ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

NDUNG'U, J.M.

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

NDUNGU, S.G.

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

In vivo comparison of susceptibility between *Bos indicus* and *Bos taurus* cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

NEVES, L.

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

NGARE, P.M.

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

NGUMI, P.N.

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

OBWOLO, M.

A macro- and light microscopical study of the pathology of *Calicophoron microbothrium* infection in experimentally infected cattle—M. MAVENYENGWA, S. MUKARATIRWA, M. OBWOLO & J. MONRAD 321–332

OHORE, O.G.

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

OLADELE, O.A.

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

OLADOKUN, A.T.

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

OLEMAPENAY, I.M.

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

OLUWAYELU, D.O.

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

OUEDRAOGO, A.G.

The phytochemical composition and *in vitro* antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (*Combretaceae*) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

PENZHORN, B.L.

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

PFUKENYI, D.M.

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

POLDER, A.

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

PROZESKY, L.

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

RANDLES, J.L.

Importation of canid rabies in a horse relocated from Zimbabwe to South Africa—C.T. SABETA & J.L. RANDLES 95–100

REYERS, F.

A canine model of normovolaemic acute anaemia—T.C. SPOTSWOOD, R.M. KIRBERGER, L.M.P.K. KOMA, F. REYERS & L. VAN DER MERWE 135–143

ROMITO, M.

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

ROTTINGHAUS, G.E.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÈ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

ROUX, C.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÈ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

SABETA, C.T.

Importation of canid rabies in a horse relocated from Zimbabwe to South Africa—C.T. SABETA & J.L. RANDLES 95–100

SALIH, D.A.

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

SAMANI, P.M.

Immunization of young chicks using graded dose of wild strain of *Eimeria tenella*—E.N. KIMBITA, P.M. SAMANI, E.N. MACHANGE & R.S. SILAYO 251–254

SCHULTZ, R.A.

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

SHARIEFF, O.E.

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

SIGAUQUE, I.

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

SILAYO, R.S.

Immunization of young chicks using graded dose of wild strain of *Eimeria tenella*—E.N. KIMBITA, P.M. SAMANI, E.N. MACHANGE & R.S. SILAYO 251–254

SMIT, F.J.

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

SPOTSWOOD, T.C.

A canine model of normovolaemic acute anaemia—T.C. SPOTSWOOD, R.M. KIRBERGER, L.M.P.K. KOMA, F. REYERS & L. VAN DER MERWE 135–143

SWART, H.

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

TARUVINGA, M.

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MACNONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

TUPPURAINEN, E.S.M.

The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques—E.S.M. TUPPURAINEN, E.H. VENTER & J.A.W. COETZER 153–164

UYS, A.C.

Ticks on crested francolins, *Francolinus sephaena*, and on the vegetation on a farm in Limpopo Province, South Africa—A.C. UYS & I.G. HORAK 339–343

VAN DEN BERG, H.

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

VAN DEN BOSSHE, P.

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

VAN DER LINDE, E.J.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

VAN DER MERWE, J.S.

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

VAN DER MERWE, L.

A canine model of normovolaemic acute anaemia—T.C. SPOTSWOOD, R.M. KIRBERGER, L.M.P.K. KOMA, F. REYERS & L. VAN DER MERWE 135–143

VAN DER WALT, S.I.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

VAN JAARSVELD, L.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

VENTER, E.H.

The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques—E.S.M. TUPPURAINEN, E.H. VENTER & J.A.W. COETZER 153–164

VILJOEN, G.J.

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

VORSTER, J.H.

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

WEBB, E.C.

Electrocardiographic surrogates of structural myocardial alterations in the Dorper sheep heart—J. KER & E.C. WEBB 273–277

WILLINGHAM, A.L.

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

WOLMARANS, C.T.

Distribution, habitats and role as intermediate host of the freshwater snail, *Bulinus forskalii*, in South Africa—K.N. DE KOCK & C.T. WOLMARANS 165–174

WOODALL, P.F.

Elephant shrews as hosts of immature ixodid ticks—L.J. FOURIE, I.G. HORAK & P.F. WOODALL 293–301

WOODFORD, M.H.

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

YOUNG, A.S.

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

ONDERSTEPOORT JOURNAL OF VETERINARY RESEARCH

Volume 72 • Numbers 1–4 • 2005

Subject index

Abamectin

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

Acari: Ixodidae

Some observations on ticks (Acari: Ixodidae) infesting sheep in River Nile Province of Northern Sudan—M. AHMED, A.M. EL HUSSEIN & A.O. EL KHIDER 239–243

Acaricides

A survey of tick control methods used by resource-poor farmers in the Qwa-Qwa area of the eastern Free State Province, South Africa—M. HLATSHWAYO & P.A. MBATI 245–249

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

Amitraz

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

Amphistomosis/amphistomes

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

Anaemia

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

A canine model of normovolaemic acute anaemia—T.C. SPOTSWOOD, R.M. KIRBERGER, L.M.P.K. KOMA, F. REYERS & L. VAN DER MERWE 135–143

Asian tapeworm

Aspects of the ecology of the Asian tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934 in yellowfish in the Vaal Dam in South Africa—A. BERTASSO & A. AVENANT-OLDEWAGE 207–217

Beak and feather disease virus

Psittacine beak and feather disease virus haemagglutinating activity using erythrocytes from African Grey parrots and Brown-headed parrots—K. KONDIAH, J. ALBERTYN & R.R. BRAGG 263–265

Birds

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

The phytochemical composition and in vitro antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (Combretaceae) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

A lectin histochemical study of the thoracic respiratory air sacs of the fowl—A.J. BEZUIDENHOUT 175–180

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Immunization of young chicks using graded dose of wild strain of *Eimeria tenella*—E.N. KIMBITA, P.M. SAMANI, E.N. MACHANGE & R.S. SILAYO 251–254

Psittacine beak and feather disease virus haemagglutinating activity using erythrocytes from African Grey parrots and Brown-headed parrots—K. KONDIAH, J. ALBERTYN & R.R. BRAGG 263–265

Ticks on crested francolins, *Francolinus sephaena*, and on the vegetation on a farm in Limpopo Province, South Africa—A.C. UYS & I.G. HORAK 339–343

Black wildebeest

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Bos indicus

In vivo comparison of susceptibility between *Bos indicus* and *Bos taurus* cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

Bos taurus

In vivo comparison of susceptibility between *Bos indicus* and *Bos taurus* cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

Bothriocephalus acheilognathi

Aspects of the ecology of the Asian tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934 in yellowfish in the Vaal Dam in South Africa—A. BERTASSO & A. AVENANT-OLDEWAGE 207–217

Bufadenolides

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

Bulinus forskalii

Distribution, habitats and role as intermediate host of the freshwater snail, *Bulinus forskalii*, in South Africa—K.N. DE KOCK & C.T. WOLMARANS 165–174

Bulinus tropicus

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

Calicophoron microbothrium

A macro- and light microscopical study of the pathology of *Calicophoron microbothrium* infection in experimentally infected cattle—M. MAVENYENGWA, S. MUKARATIRWA, M. OBWOLO & J. MONRAD 321–332

Canid rabies

Importation of canid rabies in a horse relocated from Zimbabwe to South Africa—C.T. SABETA & J.L. RANDLES 95–100

Cardenolides

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

Cardiac glycoside poisoning

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

Cardiotoxicosis

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

Cattle

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

In vivo comparison of susceptibility between *Bos indicus* and *Bos taurus* cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques—E.S.M. TUPPURAINEN, E.H. VENTER & J.A.W. COETZER 153–164

Association of trypanosomosis risk with dairy cattle production in western Kenya—G.L. MUGUNIERI & G.O. MATETE 279–284

A macro- and light microscopical study of the pathology of *Calicophoron microbothrium* infection in experimentally infected cattle—M. MAVENYENGWA, S. MUKARATIRWA, M. OBWOLO & J. MONRAD 321–332

Claviceps cyperi

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

Cypermethrin

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

***Cyperus esculentus* (nut sedge)**

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

Diminazene aceturate

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

Dogs

A preliminary study to evaluate the immune responses induced by immunization of dogs with inactivated *Ehrlichia canis* organisms—S. MAHAN, P.J. KELLY & S.M. MAHAN 119–128

Duddingtonia flagrans

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRECEK 185–187

East Coast fever

In vivo comparison of susceptibility between *Bos indicus* and *Bos taurus* cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

Ectoparasites

A survey of tick control methods used by resource-poor farmers in the Qwa-Qwa area of the eastern Free State Province, South Africa—M. HLATSHWAYO & P.A. MBATI 245–249

***Ehrlichia canis* (Ehrlichiosis)**

A preliminary study to evaluate the immune responses induced by immunization of dogs with inactivated *Ehrlichia canis* organisms—S. MAHAN, P.J. KELLY & S.M. MAHAN 119–128

Eimeria tenella

Immunization of young chicks using graded dose of wild strain of *Eimeria tenella*—E.N. KIMBITA, P.M. SAMANI, E.N. MACHANGE & R.S. SILAYO 251–254

Electrocardiography

Electrocardiographic surrogates of structural myocardial alterations in the Dorper sheep heart—J. KER & E.C. WEBB 273–277

Endometritis

Is the use of *Gunnera perpensa* extracts in endometritis related to antibacterial activity?—L.J. McGAW, R. GEHRING, L. KATSOULIS & J.N. ELOFF 129–134

Enzyme-linked immunosorbent assay

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

Ergotism

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

Fadogia homblei

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

Fish

Aspects of the ecology of the Asian tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934 in yellowfish in the Vaal Dam in South Africa—A. BERTASSO & A. AVENANT-OLDEWAGE 207–217

Fluorescence polarization immuno-assay

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

Fowl

A lectin histochemical study of the thoracic respiratory air sacs of the fowl—A.J. BEZUIDENHOUT 175–180

***Francolinus sephaena* (crested francolins)**

Ticks on crested francolins, *Francolinus sephaena*, and on the vegetation on a farm in Limpopo Province, South Africa—A.C. UYS & I.G. HORAK 339–343

Gastrodiscus aegyptiacus

Distribution, habitats and role as intermediate host of the freshwater snail, *Bulinus forskalii*, in South Africa—K.N. DE KOCK & C.T. WOLMARANS 165–174

Goats

Seroprevalence of *Toxoplasma gondii* infection in goats and sheep in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 267–272

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Gousiekte

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

Guiera senegalensis

The phytochemical composition and *in vitro* antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (*Combretaceae*) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

Gunnera perpensa

Is the use of *Gunnera perpensa* extracts in endometritis related to antibacterial activity?—L.J. McGAW, R. GEHRING, L. KATSOULIS & J.N. ELOFF 129–134

Haemonchus contortus

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRECEK 185–187

Haemophilus paragallinarum

Effects of differences in virulence of different serovars of *Haemophilus paragallinarum* on perceived vaccine efficacy —R.R. BRAGG 1–6

Heart

Electrocardiographic surrogates of structural myocardial alterations in the Dorper sheep heart—J. KER & E.C. WEBB 273–277

Heligmosomoidea

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

Helminths

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

***Hippopotamus amphibius* L (Hippopotami)**

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

Homidium chloride

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

Horse

Importation of canid rabies in a horse relocated from Zimbabwe to South Africa—C.T. SABETA & J.L. RANDLES 95–100

Hyalomma anatolicum anatolicum

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

Infectious coryza

Effects of differences in virulence of different serovars of *Haemophilus paragallinarum* on perceived vaccine efficacy —R.R. BRAGG 1–6

Isometamidium chloride

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

Kenya

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

Association of trypanosomosis risk with dairy cattle production in western Kenya—G.L. MUGUNIERI & G.O. MATETE 279–284

Kidneys

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

Labeobarbus aeneus/kimberleyensis

Aspects of the ecology of the Asian tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934 in yellowfish in the Vaal Dam in South Africa—A. BERTASSO & A. AVENANT-OLDEWAGE 207–217

Lectin binding

A lectin histochemical study of the thoracic respiratory air sacs of the fowl—A.J. BEZUIDENHOUT 175–180

Lumpy skin disease

The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques—E.S.M. TUPPURAINEN, E.H. VENTER & J.A.W. COETZER 153–164

Mammals

Small mammals as hosts of immature ixodid ticks—I.G. HORAK, L.J. FOURIE & L.E.O. BRAACK 255–261

Masaai Zebu

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

Melanoides tuberculata

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Mozambique

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

Myocardial pathology

Electrocardiographic surrogates of structural myocardial alterations in the Dorper sheep heart—J. KER & E.C. WEBB 273–277

Nematoda

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

Nematophagous fungi

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRE CEK 185–187

Nephrotoxicity

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

Nigeria

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea)

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

Nolletia gariepina

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

***Oestrus ovis/variolosus* (botflies)**

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Oriental eye-fluke

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Orma Boran

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

Ostriches

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (Struthio camelus) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Pachystigma pygmaeum

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

Parasites

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Parrots

Psittacine beak and feather disease virus haemagglutinating activity using erythrocytes from African Grey parrots and Brown-headed parrots—K. KONDIAH, J. ALBERTYN & R.R. BRAGG 263–265

Pavetta harborii

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

Pesticides

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

Philophthalmus gralli

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Pigs

Seroprevalence of *Toxoplasma gondii* infection in domestic pigs reared under different management systems in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 231–237

Plant poisoning

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

Polymerase chain reaction (PCR)

The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques—E.S.M. TUPPURAINEN, E.H. VENTER & J.A.W. COETZER 153–164

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

Premature ventricular complex

Electrocardiographic surrogates of structural myocardial alterations in the Dorper sheep heart—J. KER & E.C. WEBB 273–277

Ruminants

A study of the pathology and pathogenesis of the myocardial lesions in gousiekte, a plant-induced cardiotoxicosis of ruminants—L. PROZESKY, S.S. BASTIANELLO, N. FOURIE & R.A. SCHULTZ 219–230

Sheep

Toxicity, pathophysiology and pathology in sheep following dosing of the nephrotoxic plant *Nolletia gariepina* (DC) Mattf.—R.A. MEINTJES, C.J. BOTHA & L. PROZESKY 39–53

The role of fluorescence polarization immuno-assay in the diagnosis of plant-induced cardiac glycoside poisoning of livestock in South Africa—R.A. SCHULTZ, T.S. KELLERMAN & H. VAN DEN BERG 189–201

Some observations on ticks (Acari: Ixodidae) infesting sheep in River Nile Province of Northern Sudan—M. AHMED, A.M. EL HUSSEIN & A.O. EL KHIDER 239–243

Seroprevalence of *Toxoplasma gondii* infection in goats and sheep in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 267–272

Electrocardiographic surrogates of structural myocardial alterations in the Dorper sheep heart—J. KER & E.C. WEBB 273–277

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Snails

Distribution, habitats and role as intermediate host of the freshwater snail, *Bulinus forskalii*, in South Africa—K.N. DE KOCK & C.T. WOLMARANS 165–174

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Snotsiekte

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

South Africa

Claviceps cyperi, a new cause of severe ergotism in dairy cattle consuming maize silage and teff hay contaminated with ergotised *Cyperus esculentus* (nut sedge) on the Highveld of South Africa—T.W. NAUDÉ, C.J. BOTHA, J.H. VORSTER, C. ROUX, E.J. VAN DER LINDE, S.I. VAN DER WALT, G.E. ROTTINGHAUS, L. VAN JAARSVELD & A.N. LAWRENCE 23–37

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

Distribution, habitats and role as intermediate host of the freshwater snail, *Bulinus forskalii*, in South Africa—K.N. DE KOCK & C.T. WOLMARANS 165–174

Survey of nematophagous fungi in South Africa—D.T. DURAND, H.M. BOSHOFF, L.M. MICHAEL & R.C. KRECEK 185–187

Aspects of the ecology of the Asian tapeworm, *Bothriocephalus acheilognathi* Yamaguti, 1934 in yellowfish in the Vaal Dam in South Africa—A. BERTASSO & A. AVENANT-OLDEWAGE 207–217

A survey of tick control methods used by resource-poor farmers in the Qwa-Qwa area of the eastern Free State Province, South Africa—M. HLATSHWAYO & P.A. MBATI 245–249

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Ticks on crested francolins, *Francolinus sephaena*, and on the vegetation on a farm in Limpopo Province, South Africa—A.C. UYS & I.G. HORAK 339–343

Springbok

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Struthio camelus

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Sudan

Some observations on ticks (Acari: Ixodidae) infesting sheep in River Nile Province of Northern Sudan—M. AHMED, A.M. EL HUSSEIN & A.O. EL KHIDER 239–243

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

***Syncerus caffer* (African buffalo)**

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

Theileria annulata

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

***Theileria parva* infection (Theileriosis)**

Some preliminary observations on the susceptibility and resistance of different cattle breeds to *Theileria parva* infection—S.G. NDUNGU, P.N. NGUMI, S.K. MBOGO, T.T. DOLAN, J.J. MUTUGI & A.S. YOUNG 7–11

In vivo comparison of susceptibility between *Bos indicus* and *Bos taurus* cattle types to *Theileria parva* infection—S.G. NDUNGU, C.G.D. BROWN & T.T. DOLAN 13–22

Ticks

Some observations on ticks (Acari: Ixodidae) infesting sheep in River Nile Province of Northern Sudan—M. AHMED, A.M. EL HUSSEIN & A.O. EL KHIDER 239–243

A survey of tick control methods used by resource-poor farmers in the Qwa-Qwa area of the eastern Free State Province, South Africa—M. HLATSHWAYO & P.A. MBATI 245–249

Small mammals as hosts of immature ixodid ticks—I.G. HORAK, L.J. FOURIE & L.E.O. BRAACK 255–261

Elephant shrews as hosts of immature ixodid ticks—L.J. FOURIE, I.G. HORAK & P.F. WOODALL 293–301

Natural infection rates and transmission of *Theileria annulata* by *Hyalomma anatolicum anatolicum* ticks in the Sudan—D.A. SALIH, O.E. SHARIEFF, A.G. LAZARUS, S.M. HASSAN & A.M. EL HUSSEIN 303–307

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

Ticks on crested francolins, *Francolinus sephaena*, and on the vegetation on a farm in Limpopo Province, South Africa—A.C. UYS & I.G. HORAK 339–343

***Toxoplasma gondii* infection**

Seroprevalence of *Toxoplasma gondii* infection in domestic pigs reared under different management systems in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 231–237

Seroprevalence of *Toxoplasma gondii* infection in goats and sheep in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 267–272

Trichostrongylina

The systematic position of some Ethiopian Nippostrongylinae (Nematoda, Trichostrongylina, Heligmosomoidea) from the National Collection of Animal Helminths, Onderstepoort, South Africa—M.-C. DURETTE-DESSET & M.-C. DIGIANI 55–65

Trypanosoma congolense

The susceptibility of *Trypanosoma congolense* isolated in Zambézia Province, Mozambique, to isometamidium chloride, diminazene aceturate and homidium chloride—S. JAMAL, I. SIGAUQUE, C. MACUAMULE, L. NEVES, B.L. PENZHORN, T. MARCOTTY & P. VAN DEN BOSSCHE 333–338

Trypanosomosis

The performance of Orma Boran and Maasai Zebu crossbreeds in a trypanosomosis endemic area of Nguruman, south western Kenya—M.W. MAICHOMO, J.M. NDUNG'U, P.M. NGARE & I.M. OLEMAPENAY 87–93

Association of trypanosomosis risk with dairy cattle production in western Kenya—G.L. MUGUNIERI & G.O. MATETE 279–284

Tuberculosis

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

Uganda

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

Virus

Importation of canid rabies in a horse relocated from Zimbabwe to South Africa—C.T. SABETA & J.L. RANDLES 95–100

Serological evidence of chicken anaemia virus infection in Nigerian indigenous chickens—B.O. EMIKPE, D.O. OLUWAYELU, O.G. OHORE, O.A. OLADELE & A.T. OLADOKUN 101–103

The phytochemical composition and *in vitro* antiviral activity of decoctions from galls of *Guiera senegalensis* J.F. Gmel. (*Combretaceae*) and their relative non-toxicity for chickens—C.E. LAMIEN, A. MEDA, E. COUACY-HYMANN, A.G. OUEDRAOGO & O.G. NACOULMA 111–118

The detection of lumpy skin disease virus in samples of experimentally infected cattle using different diagnostic techniques—E.S.M. TUPPURAINEN, E.H. VENTER & J.A.W. COETZER 153–164

Psittacine beak and feather disease virus haemagglutinating activity using erythrocytes from African Grey parrots and Brown-headed parrots—K. KONDIAH, J. ALBERTYN & R.R. BRAGG 263–265

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

Widebeest

Discrimination between sheep-associated and wildebeest-associated malignant catarrhal fever virus by means of a single-tube duplex nested PCR—C.W. BREMER, H. SWART, F.A. DOBORO, B. DUNGU, M. ROMITO & G.J. VILJOEN 285–291

Wildlife

A survey of tuberculosis and other diseases in African buffalo (*Syncerus caffer*) in Queen Elizabeth National Park, Uganda—G. KALEMA-ZIKUSOKA, R.G. BENGIS, A.L. MICHEL & M.H. WOODFORD 145–151

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

Acaricide efficiency of amitraz/cypermethrin and abamectin pour-on preparations in game—J.S. VAN DER MERWE, F.J. SMIT, A.M. DURAND, L.P. KRÜGER & L.M. MICHAEL 309–314

Parasites of domestic and wild animals in South Africa. XLVI. Oestrid fly larvae of sheep, goats, springbok and black wildebeest in the Eastern Cape Province—I.G. HORAK 315–320

Zambia

Pesticide residues in adipose tissue from hippopotami (*Hippopotamus amphibius* L) living in and adjacent to the Luangwa River in Zambia—A. FLÅØYEN, A. POLDER, M. MWASE, B. ALMLI & M.M. MUSONDA 181–183

Zimbabwe

Epidemiological studies of amphistome infections in cattle in the highveld and lowveld communal grazing areas of Zimbabwe—D.M. PFUKENYI, S. MUKARATIRWA, A.L. WILLINGHAM & J. MONRAD 67–86

Importation of canid rabies in a horse relocated from Zimbabwe to South Africa—C.T. SABETA & J.L. RANDLES 95–100

First report of a field outbreak of the oriental eye-fluke, *Philophthalmus gralli*, in commercially reared ostriches (*Struthio camelus*) in Zimbabwe—S. MUKARATIRWA, THOKOZANI HOVE, Z.M. CINDZI, D.B. MAONONGA, M. TARUVINGA & ELIZABETH MATENGA 203–206

Seroprevalence of *Toxoplasma gondii* infection in domestic pigs reared under different management systems in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 231–237

Seroprevalence of *Toxoplasma gondii* infection in goats and sheep in Zimbabwe—T. HOVE, P. LIND & S. MUKARATIRWA 267–272